University of Florida Critical Care Medicine Ultrasound Curriculum (Basic Echocardiogram competencies)

Cognitive skills in recognition of clinical syndromes (based on Mayo, Paul et al. American College of Chest Physicians Statement on Competence in Critical Care Ultrasonography. Chest/135/4 April 2009)

Echocardiographic patterns: parasternal long, parasternal short, apical, subcostal
Global LV size and systolic function
Homogeneous/heterogeneous LV contraction pattern
Global RV size and systolic function
Assessment for pericardial fluid/tamponade
IVC size and respiratory variation
Basic color Doppler assessment for severe valvular regurgitation (as screening method only)
Severe hypovolemia: small, hyperdynamic ventricles; small IVC with wide respiratory variations
LV failure: Global LV systolic dysfunction; heterogeneous contractility pattern suggest of myocardial
ischemia; LV cavity dilation suggestive of chronic cardiac disease
RV failure: Acute cor pulmonale: RV dilatation and paradoxical septal motion; isolated RV dilation
suggestive of RV infarct; associated findings of dilated, no collapsible IVC
Tamponade: Pericardial effusion (regardless of size); right atrial/RV diastolic collapse; associated
findings of dilated, no collapsible IVC
Acute massive left sided valvular regurgitation: normal LV cavity size (acute valvulopathy);
normal/hyperdynamic LV systolic function (LV volume overload); massive color Doppler regurgitant
flow
Circulatory arrest during resuscitation: tamponade or acute cor pulmonale (from massive pulmonary
embolism); LV systolic function (cardiac standstill vs severely depressed vs hypderdynamic); global LV
systolic dysfunction
After successful resuscitation: heterogeneous contractility pattern suggestive of myocardial ischemia