

REVIEW ARTICLE

Julie R. Ingelfinger, M.D., *Editor*

Esophageal Carcinoma

Anil K. Rustgi, M.D., and Hashem B. El-Serag, M.D., M.P.H.

ESOPHAGEAL ADENOCARCINOMA HAS BECOME THE PREDOMINANT TYPE of esophageal cancer in North America and Europe, and gastroesophageal reflux disease (GERD) and obesity are the main risk factors. Barrett's esophagus, the recognized precursor lesion, can be detected by means of endoscopic screening, which is followed by treatment of precancerous lesions and monitoring for the development of neoplastic progression. Esophageal squamous-cell carcinoma remains the predominant esophageal cancer in Asia, Africa, and South America and among African Americans in North America. Alcohol and tobacco use are the main risk factors, and esophageal squamous dysplasia is the precursor lesion. The 5-year survival rate for patients with esophageal cancer, although generally poor, has improved during the past decade, and long-term survival is increasingly possible for patients with early or locally advanced disease. This review discusses the epidemiologic aspects and pathogenesis of these two esophageal cancers, as well as prevention and therapy, focusing on recent advances.

From the Division of Gastroenterology, Departments of Medicine and Genetics, Abramson Cancer Center, University of Pennsylvania Perelman School of Medicine, Philadelphia (A.K.R.); and the Section of Gastroenterology and Hepatology, Michael E. DeBakey Veterans Affairs Medical Center, Baylor College of Medicine, Houston (H.B.E.-S.). Address reprint requests to Dr. El-Serag at the Michael E. DeBakey VA Medical Center, 2002 Holcombe Blvd. (152), Houston, TX 77030, or at hasheme@bcm.edu.

N Engl J Med 2014;371:2499-509.

DOI: 10.1056/NEJMra1314530

Copyright © 2014 Massachusetts Medical Society.

 EPIDEMIOLOGIC ASPECTS OF ADENOCARCINOMA AND SQUAMOUS-CELL CARCINOMA

Esophageal cancer has two main subtypes — esophageal squamous-cell carcinoma and esophageal adenocarcinoma. Although squamous-cell carcinoma accounts for about 90% of cases of esophageal cancer worldwide, the incidence of and mortality rates associated with esophageal adenocarcinoma are rising and have surpassed those of esophageal squamous-cell carcinoma in several regions in North America and Europe. Esophageal carcinoma is rare in young people and increases in incidence with age, peaking in the seventh and eighth decades of life. Adenocarcinoma is three to four times as common in men as it is in women, whereas the sex distribution is more equal for squamous-cell carcinoma.

In the United States, more than 18,000 new cases of esophageal cancer and more than 15,000 deaths from esophageal cancer were expected in 2014. Over the past three decades, the rates of esophageal squamous-cell carcinoma have declined, while those of esophageal adenocarcinoma have been progressively increasing (Fig. 1).¹

 ENVIRONMENTAL RISK FACTORS

Population-based case-control and cohort studies indicate that GERD, cigarette smoking, and obesity are the main risk factors for esophageal cancer. The odds that esophageal adenocarcinoma will develop increase by a factor of five for persons with weekly GERD symptoms and by a factor of seven for persons with daily GERD symptoms, as compared with those with less frequent episodes.² The absolute risk of esophageal adenocarcinoma developing in a person 50 years of age or older is




Figure 1. Temporal Trends in Incidence Rates and Survival Rates for Esophageal Adenocarcinoma.

Blue bars denote age-adjusted incidence rates and the red line 5-year survival rates for persons with esophageal adenocarcinoma diagnosed in the United States. Data are from population-based Surveillance, Epidemiology, and End Results cancer registries.

approximately 0.04% per year.³ However, up to 40% of all patients with esophageal cancer do not report GERD symptoms.

The risk of esophageal adenocarcinoma is approximately twice as high among current smokers as it is among people who have never smoked, but smoking is a considerably stronger risk factor for esophageal squamous-cell carcinoma than for esophageal adenocarcinoma.^{4,5} In contrast, population-based studies have not shown an association between alcohol consumption and esophageal adenocarcinoma.⁶ Unlike adenocarcinoma, esophageal squamous-cell carcinoma is three to five times as likely among people who consume alcohol (three or more drinks daily), and the risk increases synergistically with tobacco smoking. High intake of red meats, fats, and processed foods is associated with an increased risk of both types of esophageal cancer, whereas high intake of fiber, fresh fruit, and vegetables is associated with a lower risk.^{7,8}

The rising incidence of esophageal adenocarcinoma has been hypothesized to be related to the increasing prevalence of GERD alone and obesity plus GERD, combined with the declining prevalence of *Helicobacter pylori* infection. Obesity is associated with a risk of esophageal adenocarcino-

ma that is increased by a factor of 2.4 to 2.8.^{9,10} Abdominal obesity in particular is associated with an increased risk of Barrett's esophagus and cancer,¹¹ possibly because increasing intragastric pressure relaxes the lower esophageal sphincter and leads to hiatal hernia, and these factors together may promote and exacerbate GERD.¹² Abdominal adiposity is more common in men, which has led to speculation that such adiposity explains some sex-related differences in cancer risk.

Populations in which *H. pylori* infection is prevalent have a reduced risk of esophageal adenocarcinoma. A meta-analysis of 15 observational studies showed that the risk of adenocarcinoma decreased by 41% among persons with *H. pylori* infection.¹³ *H. pylori* infection, which leads to gastritis, may ultimately reduce acid production through gastric atrophy, thus decreasing the exposure of the esophageal epithelium to acidic contents and reducing the risk of Barrett's esophagus^{14,15} and adenocarcinoma. However, treatment and eradication of *H. pylori* in infected patients neither causes nor exacerbates GERD in most cases.¹⁶ Overall, no consistent association between *H. pylori* and esophageal squamous-cell carcinoma has been proved.¹⁷

Esophageal adenocarcinoma has been reported in association with alendronate use.¹⁸ However, subsequent population-based studies and meta-analyses examining the association between bisphosphonate use and esophageal adenocarcinoma yielded conflicting results.¹⁹⁻²¹ Oncogenic human papillomaviruses may increase the risk of esophageal squamous-cell carcinoma, but the evidence is inconclusive.²² In addition, esophageal squamous-cell carcinoma is up to 10 times as likely to develop in patients with achalasia, an esophageal motility disorder, as it is in persons without achalasia.²³

GENETIC RISK FACTORS

Familial clustering in Barrett's esophagus and adenocarcinoma has been observed. In one genomewide, combined linkage-association analysis, germline mutations were identified in one of three candidate genes — *MSR1*, *ASCC1*, and *CTHRC1* — in 11% of patients with Barrett's esophagus or adenocarcinoma.²⁴ Mutant *MSR1* is associated with cyclin D1 overexpression, which results in more rapid cell-cycle progression.²⁵ In another

genomewide association study, susceptibility loci for Barrett's esophagus and adenocarcinoma were identified — in *CRTCI* (which encodes CREB-regulated transcription factor), *BARX1* (which encodes a protein involved in esophageal specification), and *FOXP1* (which encodes a protein involved in esophageal development).²⁶

A rare familial form of esophageal squamous-cell carcinoma — tylosis palmaris et plantaris (also called palmoplantar keratoderma), an autosomal dominant disorder characterized by hyperkeratosis of the palms and soles — has been linked to a locus on chromosome 17q21-22²⁷; missense mutations in *RHBDF2*, a gene that encodes an inactive rhomboid protease, have been identified.²⁸ Genomewide association studies have identified a number of other susceptibility loci in Chinese patients with esophageal squamous-cell carcinoma,²⁹⁻³³ which suggests that there are complex gene-environment interactions involved.

A recent study analyzed the mutational spectra from whole-exome sequencing of paired samples of tumor and normal tissue obtained from patients with esophageal adenocarcinoma and found mutations in 28 genes, 5 of which (*TP53*, *CDKN2A*, *SMAD4*, *ARID1A*, and *PIK3CA*) are relevant to the pathogenesis of adenocarcinoma.³⁴ Table S1 in the Supplementary Appendix, available with the full text of this article at NEJM.org, lists genes and molecular pathways that have been found to have alterations reported as prevalent among patients with esophageal adenocarcinoma. Lineage-specific factors, especially transcription factors, appear to be important in the development of esophageal cancers.³⁴ Likewise, whole-genome and whole-exome sequencing in Chinese patients with esophageal squamous-cell carcinoma revealed eight mutated genes — six known tumor-associated genes (*TP53*, *RBI*, *CDKN2A*, *PIK3CA*, *NOTCH1*, and *NFE2L2*) and two novel genes (*ADAM29* and *FAM135B*).³⁵

ANIMAL MODELS OF ESOPHAGEAL CANCERS

Several animal models of esophageal carcinoma are available. Surgical models in rodents, in which gastrectomy with esophagojejunal anastomosis or esophagoduodenal anastomosis is used to induce major biliary reflux, have permitted the recapitulation of Barrett's esophagus and subsequent progression to neoplasia. A mouse model in which

p120-catenin, which normally stabilizes E-cadherin at the cell membrane, is ablated solely in the esophagus results in invasive esophageal squamous-cell carcinoma.³⁶

Genetic models of esophageal adenocarcinoma also exist. For example, the esophageal-specific Epstein-Barr virus-L2 promoter, when fused to the gene encoding interleukin-1 β , induces a proinflammatory microenvironment and can induce Barrett's esophagus in mice.³⁷ Bile acid in the drinking water or crossbreeding with other mice that harbor a null allele of the *p16^{INK4a}* tumor-suppressor gene accelerates the development of esophageal adenocarcinoma. In a separate approach, the global knockout of *p63*, which is known to be critical in the proliferation of squamous epithelial stem or progenitor cells,³⁸ results in the postnatal migration of a Barrett-like cell population to the squamous-columnar junction.³⁹

In an approach that is complementary to the use of animal models, three-dimensional cell-culture “organotypic” models have been developed to elucidate pathways in the development and maintenance of Barrett's esophagus.⁴⁰⁻⁴² Barrett's esophagus may involve a switch in the fate of resident stem or progenitor cells, resulting in either reprogramming or transdifferentiation of squamous esophageal basal cells or in the migration of gastric cardia cells. After Barrett's esophagus cells emerge, dysplasia and neoplastic progression occur within the lesions, influenced by local or systemic factors (see Fig. S2 in the Supplementary Appendix). Three-dimensional organotypic culture models have also been used to study esophageal squamous-cell carcinoma.⁴³

ENDOSCOPIC SCREENING AND SURVEILLANCE

In Barrett's esophagus, which is considered the precursor of esophageal adenocarcinoma, specialized intestinal columnar epithelium replaces the normal squamous epithelium.⁴⁴ The results of large cohort studies suggest that the annual cancer risk for patients with nondysplastic Barrett's esophagus is 0.12 to 0.40%.^{45,46} Dysplasia within Barrett's esophagus lesions signals a marked increase in cancer risk — the annual risk is approximately 1% for patients with low-grade dysplasia and more than 5% for patients with high-grade dysplasia. However, 80 to 90% of cases of esophageal ad-

enocarcinoma are diagnosed in patients without known Barrett's esophagus.⁴⁷ Endoscopic screening results in detection of Barrett's esophagus in 6 to 12% of patients with prolonged GERD symptoms, most frequently white men older than 50 years of age.^{48,49} Endoscopic surveillance every 3 years is recommended for patients with known nondysplastic Barrett's esophagus.⁴⁸ Despite the absence of direct evidence from randomized trials,⁵⁰ most^{51,52} but not all⁵³ observational studies have shown that patients in whom adenocarcinoma is detected during endoscopic surveillance for Barrett's esophagus are more likely to have early-stage cancer, receive curative therapy, and survive longer than symptomatic patients in whom adenocarcinoma is detected.

In addition to leading to early detection of cancer, radiofrequency ablation of Barrett's esophagus lesions with low-grade or high-grade dysplasia results in resolution of esophageal metaplasia in up to 77% of cases and resolution of dysplasia in 86% of cases, as well as in a lower risk of progression and fewer cancers.⁵⁴ Long-term follow-up has demonstrated the durability of such effects, but continued surveillance after radiofrequency ablation is essential because of recurrences, which are mostly nondysplastic and endoscopically manageable.⁵⁵ Current guidelines do not support the use of endoscopic ablation for nondysplastic Barrett's esophagus; however, in our opinion, high-risk patients with long-segment Barrett's esophagus, severe GERD, or a family history of Barrett's esophagus or adenocarcinoma should be considered for ablative procedures. Decisions for individual patients with precancerous conditions are usually personal and unconnected to societal cost-effectiveness calculations.^{56,57} Although other tissue and blood biomarkers of Barrett's esophagus progression have been studied,⁵⁸ none have been shown to clearly outperform endoscopically detected dysplasia in predictive accuracy.

The precursor lesion for squamous-cell carcinoma is esophageal squamous dysplasia; patients with mild, moderate, or severe dysplasia have a risk of squamous-cell carcinoma that is increased by a factor of 3, 10, or 30, respectively.⁵⁹ Endoscopic screening or nonendoscopic use of balloon brush cytologic testing has been performed in some regions in China and may have merit; these techniques are also recommended for patients with achalasia or those with a history of lye ingestion resulting in stricture, although there

are no evidence-based guidelines for the treatment of these patients.

PREVENTION

PROTON-PUMP INHIBITORS

Several observational, clinic-based cohort studies have shown a significant association between treatment with proton-pump inhibitors and a decreased risk of high-grade dysplasia and adenocarcinoma in patients with Barrett's esophagus,⁶⁰⁻⁶² although limitations of these studies included possible selection bias and limited adjustment for possible confounders.⁶³ Several retrospective cohort studies have shown no reduction in the risk of esophageal adenocarcinoma among patients with GERD or Barrett's esophagus after antireflux surgery,^{64,65} which is not recommended for the sole purpose of cancer prevention.

ASPIRIN AND NSAIDS

Observational studies show a 40 to 50% reduction in the risk of esophageal adenocarcinoma and squamous-cell carcinoma with aspirin or nonsteroidal antiinflammatory drug (NSAID) treatment.⁶⁶ Given the additional cancer-reducing benefits of aspirin and NSAIDs, these medications have been recommended for general cancer chemoprevention in high-risk groups.⁶⁷ However, one randomized trial of daily celecoxib treatment did not show a reduction in cancer risk among patients with Barrett's esophagus and low-grade or high-grade dysplasia.⁶⁸ Another randomized trial showed that celecoxib did not affect the progression of esophageal squamous dysplasia.⁶⁹ Large trials examining the effects of proton-pump inhibitors and aspirin on clinical outcomes in Barrett's esophagus are ongoing.⁷⁰

STATINS

A meta-analysis of 13 studies involving humans showed a 28% reduction in the risk of esophageal adenocarcinoma among overall statin users, as compared with nonusers, and a 41% reduction in the risk of esophageal adenocarcinoma among patients with Barrett's esophagus.⁷¹ However, there was considerable inconsistency in these studies and no clear associations with dose, duration, or statin type.

The translation of other findings from epidemiologic studies into prevention recommendations — such as tobacco cessation, weight loss,⁷²

Table 1. Management of Esophageal Adenocarcinoma.

Management	Approach
Staging	Endoscopy with or without mucosal resection, computed tomography of the chest and abdomen, endoscopic ultrasonography, and positron-emission tomography
Treatment	
Mucosal tumors (stage 0 or I)*	
All tumors except T1b	Endoscopic mucosal resection (first choice) or esophagectomy with lymphadenectomy
T1b tumors	Esophagectomy with lymphadenectomy
Localized tumors (stage IIA or IIB)†	Esophagectomy preceded by neoadjuvant chemoradiotherapy (or neoadjuvant chemotherapy)
Advanced tumors (stage III or IV)	Endoscopic palliation with the use of self-expanding metal stents, with or without brachytherapy
Advanced or recurrent tumors	Two-drug or three-drug combination chemotherapy, commonly FOLFOX (infusional fluorouracil plus oxaliplatin) or XELOX (capecitabine plus oxaliplatin), or chemoradiotherapy

* Patients who are not healthy enough or are unwilling to undergo these procedures should be treated with definitive chemoradiotherapy.

† Patients who are not healthy enough or are unwilling to undergo these procedures should be treated with definitive chemoradiotherapy, especially if they have squamous-cell carcinoma.

and modification of diet to one of high fiber and low meat intake — although logical, has no clear basis. Prevention trials in China showed no benefit of nutritional supplements including vitamins and minerals in reducing the prevalence of premalignant lesions⁷³ or in reducing the incidence of or mortality associated with esophageal cancer, either in the general population⁷⁴ or among persons with esophageal squamous dysplasia.⁷⁵

CLINICAL PRESENTATION OF ESOPHAGEAL CANCER

The clinical presentation is similar between esophageal adenocarcinoma and squamous-cell carcinoma, despite differences in demographic and risk factors. The endoscopic appearance is also similar, although approximately three quarters of all adenocarcinoma lesions are found in the distal esophagus, whereas squamous-cell carcinoma is more frequent in the proximal to middle esophagus. Common clinical presentations include progressive dysphagia, weight loss, and heartburn unresponsive to medical treatment, as well as signs of blood loss; however, an increasing number of essentially asymptomatic cases are being discovered as part of screening and surveillance endoscopy. Less common symptoms include hoarseness, cough, and pneumonia related to

laryngeal nerve paralysis or invasion of the tracheobronchial tree. There is also an increased risk of synchronous and metachronous esophageal squamous-cell carcinoma in patients with head and neck squamous-cell carcinoma.

MANAGEMENT

An outline of the management of esophageal carcinoma is shown in Table 1. The management is generally similar for the two histologic types of esophageal carcinoma, except for several differences in the choice of chemotherapy or surgery. Adenocarcinoma involving the gastroesophageal junction is generally considered part of the continuum of esophageal adenocarcinoma.

STAGING

The prognosis and treatment for patients with esophageal carcinoma depend on accurate and reliable assessment of the depth of invasion and status with respect to lymph-node involvement (Fig. 2, and Table S2 in the Supplementary Appendix).⁷⁶ In the past decade, the use of endoscopic ultrasonography and positron-emission tomography (PET) has improved staging. Endoscopic ultrasonography has increased the accuracy of assessments of tumor and lymph-node status and is reported to have 70 to 80% accuracy; adding




Figure 2. Simplified Staging of Esophageal Carcinoma.

Stages 0 through IV are used to classify carcinoma characterized by different degrees of tumor invasion, lymph-node involvement, and metastasis. Stage 0 tumors are intramucosal tumors that do not invade the lamina propria. Stage I tumors invade the lamina propria without lymph-node or distant involvement. Stage II tumors extend to the muscle layer either without (IIA) or with (IIB) lymph-node involvement. Stage III tumors invade through the muscular layer and involve lymph nodes or other adjacent structures. Stage IV tumors spread to distant organs or lymph nodes. For detailed information, see Table S2 in the Supplementary Appendix.

fine-needle aspiration to endoscopic ultrasonography further improves the sensitivity of lymph-node staging.⁷⁷ Endoscopic ultrasonography is par-

ticularly helpful for staging in patients with no obvious regional or distant spread seen on imaging of the chest and abdomen; in such cases, endoscopic mucosal resection offers improved staging as well as an opportunity for cure. PET scanning identifies occult distant metastases, which are most common in the supraclavicular and retroperitoneal lymph nodes, and leads to establishment of a more advanced stage in 10 to 20% of cases.

MUCOSAL TUMORS

The introduction of endoscopic mucosal resection with or without ablation has been a major advance in treating Barrett's esophagus with high-grade dysplasia or adenocarcinoma that is limited to the epithelial portion of the mucosa (category T1a), particularly for small tumors (<2 cm in diameter) that are asymptomatic and noncircumferential. This approach is usually supplemented by endoscopic ablation of the remaining Barrett's esophagus lesions (Fig. 3). The risk of lymph-node metastasis is correlated with the depth of tumor invasion; the risk is close to zero among patients with Barrett's esophagus who have high-grade dysplasia and is only 1 to 2% among patients with stage I tumors.⁷⁸ There are no data from randomized trials comparing endoscopic therapies with surgical approaches,⁷⁹ but several observational studies have suggested that cure and survival rates associated with endoscopic treatments are equivalent to the rates with surgical resection.⁸⁰ Endoscopic therapy should be considered as first-line therapy for patients with stage 0 or I esophageal adenocarcinoma who do not have contraindications or major coexisting conditions.⁸¹ In patients with category T1b tumors that have penetrated the muscularis mucosae and entered the submucosa, the risk of lymph-node spread is as high as 20%, and radical esophagectomy may be the preferred method of treatment, although some treatment centers have expanded the indications for endoscopic therapy to include low-risk submucosal tumors.⁸²

LOCALLY ADVANCED TUMORS

Locally advanced tumors, defined as category T3N1, are best treated with esophagectomy. Although cure through definitive chemoradiotherapy alone has been reported, especially in patients with squamous-cell carcinoma,⁸³ that approach is not supported by evidence from randomized, controlled trials and should be restricted to pa-




Figure 3. Endoscopic Images.

Panel A shows intramucosal adenocarcinoma in the background of Barrett's esophagus at the lower end of the esophagus. Panel B is an endoscopic ultrasound image showing esophageal mucosa thickening (orange arrow) and intact submucosa (blue arrow). Panel C shows endoscopic mucosal resection with the use of a suction cap, followed by band application and resection with a thermal snare. Panel D is the view after endoscopic mucosal resection, showing exposed clean submucosa. Panel E is the view after radiofrequency ablation of the remaining Barrett's esophagus, showing superficial circumferential ulceration and denuding of the Barrett's esophagus. Panel F shows the healed esophagus approximately 3 months later. Follow-up biopsy results showed normal squamous epithelium with no metaplasia, dysplasia, or cancer.

tients whose condition is declining or who are not healthy enough to undergo esophagectomy. Unfortunately, esophagectomy alone is associated with a high rate of recurrence and low 5-year survival rates (5 to 34%). The main advance in treating patients who undergo esophagectomy has been the adoption of neoadjuvant treatment. Randomized, controlled trials have shown a survival benefit with neoadjuvant chemoradiotherapy or chemotherapy, as compared with esophagectomy alone, in both types of esophageal carcinoma.⁸³⁻⁸⁵ Chemoradiotherapy with carboplatin and paclitaxel⁸⁴ or cisplatin and fluorouracil⁸⁵ is becoming the standard treatment in the United States⁸⁶; in Europe, neoadjuvant chemotherapy alone is the preferred approach. There may be a small advantage to neoadjuvant chemoradiotherapy over chemotherapy alone. A meta-analysis showed a pooled hazard ratio for death from any cause of 0.78 (95% confidence interval [CI], 0.70 to 0.88) for neoadjuvant chemoradiotherapy and 0.87 (95% CI, 0.79 to 0.96) for neoadjuvant chemotherapy, with

a greater benefit in adenocarcinoma than in squamous-cell carcinoma.⁸⁷ Chemotherapy before and after resection may have a small additional benefit in patients with squamous-cell carcinoma.⁸⁸ There is no reliable test apart from histologic examination of the resected specimen to confirm the response to neoadjuvant therapy, and esophagectomy therefore remains necessary. Patients with adenocarcinoma who have residual, node-positive, completely resected disease after neoadjuvant therapy have poor outcomes, and the benefits of adjuvant chemotherapy or chemoradiotherapy are unclear in such patients. However, adjuvant chemotherapy is used for node-positive patients with squamous-cell carcinoma; this approach has been shown to have benefits in several randomized trials in Japan.⁸⁹

Surgical outcomes appear to be better in high-volume centers and with experienced surgeons, a benefit apparently related to the incidence and management of postoperative complications.⁹⁰ Two-field lymphadenectomy in the abdomen and tho-

rax is standard practice in most centers in Europe and North America, and additional dissection of lymph nodes of the neck is performed in some countries where squamous-cell carcinoma is common. Minimally invasive esophagectomy may have equivalent safety and a similar rate of complications, but it is technically challenging, even when performed in specialized centers.⁹¹

ADVANCED TUMORS

Obstructive symptoms related to unresectable disease can be palliated with endoscopic esophageal stenting (see Fig. S2 in the Supplementary Appendix) or high-dose intraluminal brachytherapy.⁹² Endoscopic placement of self-expanding metal stents has become the first-line palliative option for dysphagia. Randomized, controlled trials have shown higher symptomatic relief, as well as less need for reintervention due to complications, with self-expanding metal stents than with locoregional treatment. The addition of high-dose brachytherapy to stenting may result in a modest prolongation of survival.⁹³ Other methods of treatment, such as endoscopic dilation or ablation, placement of plastic stents, bypass surgery, or chemoradiotherapy, are not recommended because of their low efficacy and high rates of complications.

Palliative chemotherapy for the prolongation of survival is also commonly used to treat patients with unresectable, metastatic, or recurrent disease. Treatment with cisplatin or oxaliplatin combined with either infusional fluorouracil or capecitabine achieves response rates of 35 to 45% and a few months of prolonged survival, especially among patients with squamous-cell carcinoma. The addition of a third drug may increase response rates by an additional 5 to 10 percentage points but is associated with higher toxicity. Second-line chemotherapy remains investigational. The use of gefitinib as a second-line treatment for unselected patients does not improve overall survival.⁹⁴ Promising results are emerging with docetaxel (which interferes with cell division by stabilizing microtubules)⁹⁵ and ramucirumab (which targets vascular endothelial growth factor receptor 2). Trastuzumab, a biologic agent in which ERBB2 is amplified, confers a modest 2.7-month increase in overall survival and a 1.7-month increase in progression-free survival among patients with advanced esophageal adenocarcinomas.⁹⁶

PROGNOSIS

The overall 5-year survival rate for patients with esophageal adenocarcinoma in the United States is approximately 17%, which is slightly higher than the rate for patients with squamous-cell carcinoma (Fig. 1). There has been a progressive improvement in overall survival and a marked improvement in progression-free survival among patients who undergo surgical resection. In spite of the fact that the ability to detect early-stage esophageal adenocarcinoma has improved, most tumors are found when regional metastasis (in 30% of cases) or distant metastasis (in 40% of cases) has already occurred, at which point the 5-year survival rate declines from 39% in cases of localized disease to 4% in cases with distant metastasis. Furthermore, 60 to 70% of patients with esophageal cancer have not been receiving guideline-concordant treatment. The management of esophageal cancer appears to be improved by discussion with a multidisciplinary tumor board.

SUMMARY

The main risk factors for esophageal adenocarcinoma are GERD, obesity, and cigarette smoking; *H. pylori* infection is associated with a reduced risk. Cigarette smoking and alcohol consumption constitute the main risk factors for esophageal squamous-cell carcinoma. Endoscopic screening allows for the identification of Barrett's esophagus, which in turn allows for periodic surveillance of Barrett's esophagus for the detection of dysplasia and early-stage esophageal adenocarcinoma. Endoscopic ablative therapy has been shown to be efficacious for the treatment of dysplasia and may have an important role in the treatment of intramucosal adenocarcinoma. Identification of certain genomic regions and genes might provide insights into the underlying pathogenesis of esophageal cancer and may have translational implications for biomarker identification and development of novel therapies.

The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs or the U.S. government.

No potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

REFERENCES

- Hur C, Miller M, Kong CY, et al. Trends in esophageal adenocarcinoma incidence and mortality. *Cancer* 2013;119:1149-59.
- Rubenstein JH, Taylor JB. Meta-analysis: the association of oesophageal adenocarcinoma with symptoms of gastro-oesophageal reflux. *Aliment Pharmacol Ther* 2010;32:1222-7.
- Shaheen N, Ransohoff DF. Gastroesophageal reflux, Barrett esophagus, and esophageal cancer: scientific review. *JAMA* 2002;287:1972-81.
- Cook MB, Kamangar F, Whitman DC, et al. Cigarette smoking and adenocarcinomas of the esophagus and esophagogastric junction: a pooled analysis from the international BEACON consortium. *J Natl Cancer Inst* 2010;102:1344-53.
- Tramacere I, La Vecchia C, Negri E. Tobacco smoking and alcohol drinking and gastric cardia adenocarcinoma: a meta-analysis. *Epidemiology* 2011;22:344-9.
- Tramacere I, Pelucchi C, Bagnardi V, et al. A meta-analysis on alcohol drinking and esophageal and gastric cardia adenocarcinoma risk. *Ann Oncol* 2012;23:287-97.
- Kubo A, Corley DA, Jensen CD, Kaur R. Dietary factors and the risks of oesophageal adenocarcinoma and Barrett's oesophagus. *Nutr Res Rev* 2010;23:230-46.
- Coleman HG, Murray LJ, Hicks B, et al. Dietary fiber and the risk of precancerous lesions and cancer of the esophagus: a systematic review and meta-analysis. *Nutr Rev* 2013;71:474-82.
- Kubo A, Corley DA. Body mass index and adenocarcinomas of the esophagus or gastric cardia: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:872-8.
- Hampel H, Abraham NS, El-Serag HB. Meta-analysis: obesity and the risk for gastroesophageal reflux disease and its complications. *Ann Intern Med* 2005;143:199-211.
- Singh S, Sharma AN, Murad MH, et al. Central adiposity is associated with increased risk of esophageal inflammation, metaplasia, and adenocarcinoma: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2013;11:1399-412.
- El-Serag H. Role of obesity in GORD-related disorders. *Gut* 2008;57:281-4.
- Xie FJ, Zhang YP, Zheng QQ, et al. Helicobacter pylori infection and esophageal cancer risk: an updated meta-analysis. *World J Gastroenterol* 2013;19:6098-107.
- Fischbach LA, Graham DY, Kramer JR, et al. Association between Helicobacter pylori and Barrett's esophagus: a case-control study. *Am J Gastroenterol* 2014;109:357-68.
- Rubenstein JH, Inadomi JM, Scheiman J, et al. Association between Helicobacter pylori and Barrett's esophagus, erosive esophagitis, and gastroesophageal reflux symptoms. *Clin Gastroenterol Hepatol* 2014;12:239-45.
- Malfertheiner P, Megraud F, O'Morain CA, et al. Management of Helicobacter pylori infection — the Maastricht IV/ Florence Consensus Report. *Gut* 2012;61:646-64.
- Rokkas T, Pistiolos D, Sechopoulos P, Robotis I, Margantinis G. Relationship between Helicobacter pylori infection and esophageal neoplasia: a meta-analysis. *Clin Gastroenterol Hepatol* 2007;5:1413-7.
- Wysowski DK. Reports of esophageal cancer with oral bisphosphonate use. *N Engl J Med* 2009;360:89-90.
- Cardwell CR, Abnet CC, Cantwell MM, Murray LJ. Exposure to oral bisphosphonates and risk of esophageal cancer. *JAMA* 2010;304:657-63.
- Green J, Czanner G, Reeves G, Watson J, Wise L, Beral V. Oral bisphosphonates and risk of cancer of oesophagus, stomach, and colorectum: case-control analysis within a UK primary care cohort. *BMJ* 2010;341:c4444.
- Ho YF, Lin JT, Wu CY. Oral bisphosphonates and risk of esophageal cancer: a dose-intensity analysis in a nationwide population. *Cancer Epidemiol Biomarkers Prev* 2012;21:993-5.
- Hardefeldt HA, Cox MR, Eslick GD. Association between human papillomavirus (HPV) and oesophageal squamous cell carcinoma: a meta-analysis. *Epidemiol Infect* 2014;142:1119-37.
- Zendehdel K, Nyren O, Edberg A, Ye W. Risk of esophageal adenocarcinoma in achalasia patients, a retrospective cohort study in Sweden. *Am J Gastroenterol* 2011;106:57-61.
- Orloff M, Peterson C, He X, et al. Germline mutations in MSR1, ASCCL1, and CTHRC1 in patients with Barrett esophagus and esophageal adenocarcinoma. *JAMA* 2011;306:410-9.
- Shan J, Zhao W, Gu W. Suppression of cancer cell growth by promoting cyclin D1 degradation. *Mol Cell* 2009;36:469-76.
- Levine DM, Ek WE, Zhang R, et al. A genome-wide association study identifies new susceptibility loci for esophageal adenocarcinoma and Barrett's esophagus. *Nat Genet* 2013;45:1487-93.
- Reis A, Küster W, Eckardt R, Sperling K. Mapping of a gene for epidermolytic palmoplantar keratoderma to the region of the acidic keratin gene cluster at 17q12-q21. *Hum Genet* 1992;90:113-6.
- Blaydon DC, Etheridge SL, Risk JM, et al. RHBDF2 mutations are associated with tylosis, a familial esophageal cancer syndrome. *Am J Hum Genet* 2012;90:340-6.
- Wu C, Li D, Jia W, et al. Genome-wide association study identifies common variants in SLC39A6 associated with length of survival in esophageal squamous-cell carcinoma. *Nat Genet* 2013;45:632-8.
- Wu C, Kraft P, Zhai K, et al. Genome-wide association analyses of esophageal squamous cell carcinoma in Chinese identify multiple susceptibility loci and gene-environment interactions. *Nat Genet* 2012;44:1090-7.
- Wu C, Hu Z, He Z, et al. Genome-wide association study identifies three new susceptibility loci for esophageal squamous-cell carcinoma in Chinese populations. *Nat Genet* 2011;43:679-84.
- Wang LD, Zhou FY, Li XM, et al. Genome-wide association study of esophageal squamous cell carcinoma in Chinese subjects identifies susceptibility loci at PLCE1 and C20orf54. *Nat Genet* 2010;42:759-63.
- Abnet CC, Freedman ND, Hu N, et al. A shared susceptibility locus in PLCE1 at 10q23 for gastric adenocarcinoma and esophageal squamous cell carcinoma. *Nat Genet* 2010;42:764-7.
- Dulak AM, Stojanov P, Peng S, et al. Exome and whole-genome sequencing of esophageal adenocarcinoma identifies recurrent driver events and mutational complexity. *Nat Genet* 2013;45:478-86.
- Song Y, Li L, Ou Y, et al. Identification of genomic alterations in oesophageal squamous cell cancer. *Nature* 2014;509:91-5.
- Stairs DB, Bayne LJ, Rhoades B, et al. Deletion of p120-catenin results in a tumor microenvironment with inflammation and cancer that establishes it as a tumor suppressor gene. *Cancer Cell* 2011;19:470-83.
- Quante M, Bhagat G, Abrams JA, et al. Bile acid and inflammation activate gastric cardia stem cells in a mouse model of Barrett-like metaplasia. *Cancer Cell* 2012;21:36-51.
- Senoo M, Pinto F, Crum CP, McKeon F. p63 is essential for the proliferative potential of stem cells in stratified epithelia. *Cell* 2007;129:523-36.
- Wang X, Ouyang H, Yamamoto Y, et al. Residual embryonic cells as precursors of a Barrett's-like metaplasia. *Cell* 2011;145:1023-35.
- Stairs DB, Nakagawa H, Klein-Szanto A, et al. Cdx1 and c-Myc foster the initiation of transdifferentiation of the normal esophageal squamous epithelium toward Barrett's esophagus. *PLoS One* 2008;3(10):e3534.
- Kosoff RE, Gardiner KL, Merlo LM, Pavlov K, Rustgi AK, Maley CC. Development and characterization of an organotypic model of Barrett's esophagus. *J Cell Physiol* 2012;227:2654-9.

42. Bass AJ, Wang TC. An inflammatory situation: SOX2 and STAT3 cooperate in squamous cell carcinoma initiation. *Cell Stem Cell* 2013;12:266-8.
43. Okawa T, Michaylira CZ, Kalabis J, et al. The functional interplay between EGFR overexpression, hTERT activation, and p53 mutation in esophageal epithelial cells with activation of stromal fibroblasts induces tumor development, invasion, and differentiation. *Genes Dev* 2007;21:2788-803.
44. Spechler SJ, Souza RF. Barrett's esophagus. *N Engl J Med* 2014;371:836-45.
45. Hvid-Jensen F, Pedersen L, Drewes AM, Sørensen HT, Funch-Jensen P. Incidence of adenocarcinoma among patients with Barrett's esophagus. *N Engl J Med* 2011;365:1375-83.
46. de Jonge PJ, van Blankenstein M, Looman CW, Casparie MK, Meijer GA, Kuipers EJ. Risk of malignant progression in patients with Barrett's oesophagus: a Dutch nationwide cohort study. *Gut* 2010;59:1030-6.
47. Dulai GS, Guha S, Kahn KL, Gornbein J, Weinstein WM. Preoperative prevalence of Barrett's esophagus in esophageal adenocarcinoma: a systematic review. *Gastroenterology* 2002;122:26-33.
48. Spechler SJ, Sharma P, Souza RF, Inadomi JM, Shaheen NJ. American Gastroenterological Association technical review on the management of Barrett's esophagus. *Gastroenterology* 2011;140(3):e18-e52.
49. Sharma P. Barrett's esophagus. *N Engl J Med* 2009;361:2548-56. [Erratum, *N Engl J Med* 2010;362:1450.]
50. Yang S, Wu S, Huang Y, et al. Screening for oesophageal cancer. *Cochrane Database Syst Rev* 2012;12:CD007883.
51. Cooper GS, Kou TD, Chak A. Receipt of previous diagnoses and endoscopy and outcome from esophageal adenocarcinoma: a population-based study with temporal trends. *Am J Gastroenterol* 2009;104:1356-62.
52. Kearney DJ, Crump C, Maynard C, Boyko EJ. A case-control study of endoscopy and mortality from adenocarcinoma of the esophagus or gastric cardia in persons with GERD. *Gastrointest Endosc* 2003;57:823-9.
53. Corley DA, Levin TR, Habel LA, Weiss NS, Buffler PA. Surveillance and survival in Barrett's adenocarcinomas: a population-based study. *Gastroenterology* 2002;122:633-40.
54. Shaheen NJ, Sharma P, Overholt BF, et al. Radiofrequency ablation in Barrett's esophagus with dysplasia. *N Engl J Med* 2009;360:2277-88.
55. Orman ES, Li N, Shaheen NJ. Efficacy and durability of radiofrequency ablation for Barrett's esophagus: systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2013;11:1245-55.
56. El-Serag HB, Graham DY. Routine polypectomy for colorectal polyps and ablation for Barrett's esophagus are intellectually the same. *Gastroenterology* 2011;140:386-8.
57. El-Serag HB, Naik AD. Surveillance in Barrett's esophagus: lessons from behavioral economics. *Gastroenterology* 2009;137:763-5.
58. Bird-Lieberman EL, Dunn JM, Coleman HG, et al. Population-based study reveals new risk-stratification biomarker panel for Barrett's esophagus. *Gastroenterology* 2012;143:927-35.
59. Wang GQ, Abnet CC, Shen Q, et al. Histological precursors of oesophageal squamous cell carcinoma: results from a 13 year prospective follow up study in a high risk population. *Gut* 2005;54:187-92.
60. Nguyen DM, El-Serag HB, Henderson L, Stein D, Bhattacharyya A, Sampliner RE. Medication usage and the risk of neoplasia in patients with Barrett's esophagus. *Clin Gastroenterol Hepatol* 2009;7:1299-304.
61. Hillman LC, Chiragakis L, Shadbolt B, Kaye GL, Clarke AC. Proton-pump inhibitor therapy and the development of dysplasia in patients with Barrett's esophagus. *Med J Aust* 2004;180:387-91.
62. Kastelein F, Spaander MC, Steyerberg EW, et al. Proton pump inhibitors reduce the risk of neoplastic progression in patients with Barrett's esophagus. *Clin Gastroenterol Hepatol* 2013;11:382-8.
63. Singh S, Garg SK, Singh PP, Iyer PG, El-Serag HB. Acid-suppressive medications and risk of oesophageal adenocarcinoma in patients with Barrett's oesophagus: a systematic review and meta-analysis. *Gut* 2014;63:1229-37.
64. Corey KE, Schmitz SM, Shaheen NJ. Does a surgical antireflux procedure decrease the incidence of esophageal adenocarcinoma in Barrett's esophagus? A meta-analysis. *Am J Gastroenterol* 2003;98:2390-4.
65. Tran T, Spechler SJ, Richardson P, El-Serag HB. Fundoplication and the risk of esophageal cancer in gastroesophageal reflux disease: a Veterans Affairs cohort study. *Am J Gastroenterol* 2005;100:1002-8.
66. Liao LM, Vaughan TL, Corley DA, et al. Nonsteroidal anti-inflammatory drug use reduces risk of adenocarcinomas of the esophagus and esophagogastric junction in a pooled analysis. *Gastroenterology* 2012;142:442-52.
67. Chan AT, Detering E. An emerging role for anti-inflammatory agents for chemoprevention. *Recent Results Cancer Res* 2013;191:1-5.
68. Heath EL, Canto MI, Piantadosi S, et al. Secondary chemoprevention of Barrett's esophagus with celecoxib: results of a randomized trial. *J Natl Cancer Inst* 2007;99:545-57.
69. Limburg PJ, Wei W, Ahnen DJ, et al. Randomized, placebo-controlled, esophageal squamous cell cancer chemoprevention trial of selenomethionine and celecoxib. *Gastroenterology* 2005;129:863-73.
70. Das D, Chilton AP, Jankowski JA. Chemoprevention of oesophageal cancer and the AsPECT trial. *Recent Results Cancer Res* 2009;181:161-9.
71. Singh S, Singh AG, Singh PP, Murad MH, Iyer PG. Statins are associated with reduced risk of esophageal cancer, particularly in patients with Barrett's esophagus: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2013;11:620-9.
72. Scozzari G, Trapani R, Toppino M, Morino M. Esophagogastric cancer after bariatric surgery: systematic review of the literature. *Surg Obes Relat Dis* 2013;9:133-42.
73. Muñoz N, Wahrendorf J, Bang LJ, et al. No effect of riboflavine, retinol, and zinc on prevalence of precancerous lesions of oesophagus: randomised double-blind intervention study in high-risk population of China. *Lancet* 1985;2:111-4.
74. Blot WJ, Li JY, Taylor PR, et al. Nutrition intervention trials in Linxian, China: supplementation with specific vitamin/mineral combinations, cancer incidence, and disease-specific mortality in the general population. *J Natl Cancer Inst* 1993;85:1483-92.
75. Li JY, Taylor PR, Li B, et al. Nutrition intervention trials in Linxian, China: multiple vitamin/mineral supplementation, cancer incidence, and disease-specific mortality among adults with esophageal dysplasia. *J Natl Cancer Inst* 1993;85:1492-8.
76. Pennathur A, Gibson MK, Jobe BA, Luketich JD. Oesophageal carcinoma. *Lancet* 2013;381:400-12.
77. van Vliet EP, Heijnenbroek-Kal MH, Hunink MG, Kuipers EJ, Siersema PD. Staging investigations for oesophageal cancer: a meta-analysis. *Br J Cancer* 2008;98:547-57.
78. Dunbar KB, Spechler SJ. The risk of lymph-node metastases in patients with high-grade dysplasia or intramucosal carcinoma in Barrett's esophagus: a systematic review. *Am J Gastroenterol* 2012;107:850-62.
79. Bennett C, Green S, Decaestecker J, et al. Surgery versus radical endotherapies for early cancer and high-grade dysplasia in Barrett's oesophagus. *Cochrane Database Syst Rev* 2012;11:CD007334.
80. Ngamruengphong S, Wolfson HC, Wallace MB. Survival of patients with superficial esophageal adenocarcinoma after endoscopic treatment vs surgery. *Clin Gastroenterol Hepatol* 2013;11:1424-9.
81. Fitzgerald RC, di Pietro M, Ragunath K, et al. British Society of Gastroenterology guidelines on the diagnosis and management of Barrett's oesophagus. *Gut* 2014;63:7-42.
82. Pech O, Behrens A, May A, et al. Long-term results and risk factor analysis for

- recurrence after curative endoscopic therapy in 349 patients with high-grade intraepithelial neoplasia and mucosal adenocarcinoma in Barrett's oesophagus. *Gut* 2008;57:1200-6.
83. Morgan MA, Lewis WG, Casbard A, et al. Stage-for-stage comparison of definitive chemoradiotherapy, surgery alone and neoadjuvant chemotherapy for oesophageal carcinoma. *Br J Surg* 2009;96:1300-7.
84. van Hagen P, Hulshof MC, van Lanschot JJ, et al. Preoperative chemoradiotherapy for esophageal or junctional cancer. *N Engl J Med* 2012;366:2074-84.
85. Tepper J, Krasna MJ, Niedzwiecki D, et al. Phase III trial of trimodality therapy with cisplatin, fluorouracil, radiotherapy, and surgery compared with surgery alone for esophageal cancer: CALGB 9781. *J Clin Oncol* 2008;26:1086-92.
86. Malthaner R, Wong RK, Spithoff K. Preoperative or postoperative therapy for resectable oesophageal cancer: an updated practice guideline. *Clin Oncol (R Coll Radiol)* 2010;22:250-6.
87. Sjoquist KM, Burmeister BH, Smithers BM, et al. Survival after neoadjuvant chemotherapy or chemoradiotherapy for resectable oesophageal carcinoma: an updated meta-analysis. *Lancet Oncol* 2011;12:681-92.
88. Cunningham D, Allum WH, Stenning SP, et al. Perioperative chemotherapy versus surgery alone for resectable gastroesophageal cancer. *N Engl J Med* 2006;355:11-20.
89. Ando N, Iizuka T, Ide H, et al. Surgery plus chemotherapy compared with surgery alone for localized squamous cell carcinoma of the thoracic esophagus: a Japan Clinical Oncology Group Study — JCOG9204. *J Clin Oncol* 2003;21:4592-6.
90. Wouters MW, Gooiker GA, van Sandick JW, Tollenaar RA. The volume-outcome relation in the surgical treatment of esophageal cancer: a systematic review and meta-analysis. *Cancer* 2012;118:1754-63.
91. Herbella FA, Patti MG. Minimally invasive esophagectomy. *World J Gastroenterol* 2010;16:3811-5.
92. Sreedharan A, Harris K, Crellin A, Forman D, Everett SM. Interventions for dysphagia in oesophageal cancer. *Cochrane Database Syst Rev* 2009;4:CD005048.
93. Zhu HD, Guo JH, Mao AW, et al. Conventional stents versus stents loaded with (125)iodine seeds for the treatment of unresectable oesophageal cancer: a multicentre, randomised phase 3 trial. *Lancet Oncol* 2014;15:612-9.
94. Dutton SJ, Ferry DR, Blazeby JM, et al. Gefitinib for oesophageal cancer progressing after chemotherapy (COG): a phase 3, multicentre, double-blind, placebo-controlled randomised trial. *Lancet Oncol* 2014;15:894-904.
95. Ford HE, Marshall A, Bridgewater JA, et al. Docetaxel versus active symptom control for refractory oesophagogastric adenocarcinoma (COUGAR-02): an open-label, phase 3 randomised controlled trial. *Lancet Oncol* 2014;15:78-86.
96. Bang YJ, Van Cutsem E, Feyereislova A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet* 2010;376:687-97. [Erratum, *Lancet* 2010;376:1302.]

Copyright © 2014 Massachusetts Medical Society.

RECEIVE IMMEDIATE NOTIFICATION WHEN AN ARTICLE
IS PUBLISHED ONLINE FIRST

To be notified by e-mail when *Journal* articles
are published Online First, sign up at NEJM.org.